

# Influence of organic matter on the secondary urban effluents disinfection by UV/H2O2

## Ana R. Neves\*, Jacqueline A. Malvestiti, Renato F. Dantas.

## Abstract

The variation in the composition and the reactivity of the effluent can delay the inactivation of microorganisms and the degradation of organic contaminants due to the presence of reactive material that competes to react with the oxidants. In the case of advanced oxidative processes, the radicals formed, especially HO. can be affected by the organic matter reactivity (EfOM). This project proposed the study of the influence of organic matter on the efficiency of UV/H2O2 disinfection through the measurement of total coliforms and E. coli. The experiments were carried out with Milig-water, samples from the secondary effluent treatment plant of School of Technology, University of Campinas - UNICAMP and from Águas da Serra - wastewater treatment plant (WWTP), Limeira-SP. The relation between the organic matter of the effluent and the disinfection process was studied from the coliform concentration data compared to the COD of the effluent. For samples from the Águas da Serra effluent, which presented COD of 79 mg/L, the disinfection was extremely efficient with different concentrations of peroxide and there was no indication of the presence of coliforms from the first 10 minutes of treatment. The samples from Faculdade de Tecnologia treatment plant presented COD of 96,2 mg/L and did not show total disinfection in the first 10 minutes with the peroxide concentration of 45mg/L.

Key words: Advanced oxidation processes, disinfection, UV/H2O2

#### Introduction

Nowadays there is a continuously increasing worldwide concern for development of alternative water reuse technologies, mainly focused on agriculture and industry. In this context, Advanced Oxidation Processes (AOPs) are considered a highly competitive water treatment technology for the removal of those organic pollutants not treatable by conventional techniques due to their high chemical stability and/or low. Numerous studies report that the UV/H<sub>2</sub>O<sub>2</sub>, is effective for the destruction of various pollutants however there are very few available reports on inactivation of microorganisms via OH radical oxidation (Mamane et al., 2007). Although the hydroxyl radical (HO-) is the main oxidizing agent in these processes (oxidative power: 2.80 eV) (Ikehata et al., 2006).

The objective of this research was to study the influence of organic matter and carbonate on the disinfection process of UV/H<sub>2</sub>O<sub>2</sub> through the inactivation of total coliforms and E. coli.

## **Results and Discussion**

The average of Águas da Serra treatment plant COD was 79 mg/L. For these samples, the total coliform and E. coli were totally inactivated after 10 minutes of treatment. Chart 1. Águas da Serra treatment plant (30mg H<sub>2</sub>O<sub>2</sub>/L)





The effluent from School of Technology, University of Campinas presented COD of 96.2 mg/L and the disinfection rate of the following graph for 45 mg of H<sub>2</sub>O<sub>2</sub>/L.

Chart 2. Faculdade de Tecnologia treatment plant (45mg H2O2/L)



#### Conclusions

It can be concluded that the disinfection rate in the first 10 minutes of treatment is higher for the effluent of the Aguas da Serra treatment plant than for the effluent from the School of Technology, University of Campinas.

#### Acknowledgement

I thank Jacqueline Malvestiti and Raissa Barbacena for their assistance. I also thank PIBIC for the funding.

Mamane, H.; Shemer, H.; Linden, K. G. Inactivation of E. coli, B. subtilis spores, and MS2, T4, and T7 phage using UV/H2O2 advanced oxidation. Journal of Hazardous Materials 146 (2007), p. 479–486; <sup>1</sup> Curtis, M. D.; Shiu, K.; Butler, W. M. e Huffmann, J. C. J. Am. Chem. Soc. 1986, 108, 3335.

Ikehata, K., Jodeiri Naghashkar, N., Gamal El-Din, M., 2006. Degradation of aqueous pharmaceuticals by ozonation and advanced oxidation processes: a review. Ozone Sci. Eng. 28 (6), 353-414.

